

Odderon and Pomeron from the Vacuum Correlator Method

A.B.Kaidalov, Yu.A.Simonov

State Research Center

Institute of Theoretical and Experimental Physics,
Moscow, 117218 Russia

Abstract

Glueball masses with $J \leq 7$ are computed both for $C = +1$ and $C = -1$ using the string Hamiltonian derived in the framework of the Vacuum Correlator Method. No fitting parameters are used, and masses are expressed in terms of string tension σ and effective value of α_s .

We extend the calculations done for $J \leq 3$ using the same Hamiltonian, which provided glueball masses in good agreement with existing lattice data, to higher mass states. It is shown that 3^{--} , 5^{--} and 7^{--} states lie on the odderon trajectories with the intercept around or below 0.14. Another odderon trajectory with $3g$ glueballs of Y -shape, corresponds to 11% higher masses and low intercept. These findings are in agreement with recent experimental data, setting limits on the odderon contribution to the exclusive γp reactions.

1 Introduction

There was a renewal of interest in glueballs recently, mostly connected to pomeron and odderon trajectories [1]-[6]. High-spin glueballs, e.g. 4^{++} , 6^{++} were calculated on the lattice [1, 2], and the problem of odderon attracted much attention [3, 4, 5]. Search for odderon exchange at HERA [6, 7] in the reactions $\gamma p \rightarrow \pi^0 p$, $\gamma p \rightarrow \pi^0 \pi^0 p$, ... has not provided any indications for an

existence of these processes at high energies¹ and has lead to rather strong limits on the corresponding cross sections. This allows to put limits on the intercept of the odderon in the models which predict couplings of the odderon with hadrons [4]. Study of this problem allows to distinguish between the models and seriously question some of them. Therefore it seems to be necessary to make a more detailed analysis of our theoretical calculations [8] and to compare additional high-spin glueball masses.

A theoretical study of glueballs in QCD was started in [9]-[12] and is closely related to the problem of the pomeron, i.e. leading Regge pole, which determines the asymptotic behavior of scattering amplitudes at very high energies. It is usually assumed that the pomeron in QCD is mostly gluonic object [13] and glueball resonances with vacuum quantum numbers and spins belong to this trajectory. Another interesting hypothetical Regge singularity is the "odderon", which has negative signature and C parity and can be built out of at least 3 gluons. Most studies of the pomeron and odderon singularities in QCD are based on applications of the perturbation theory [14].

Our method is based on the QCD path integral formalism, where all dynamics is encoded in field correlators – the so-called Field Correlator Method (FCM) [15] (for a review see [16]) starting from that one can derive in the limit of small gluon correlation length λ the relativistic Hamiltonian [17], which effectively describes the fundamental (adjoint) string with quarks (gluons) at its ends. In this simple limit, $\lambda \rightarrow 0$, Hamiltonian is local but nonlinear in $\mathbf{p}^2, \hat{L}^2$, which reflects complicated dynamics of relativistic rotating string (similar results are obtained in [18]).

One should stress, that nonperturbative (NP) approach started in [19] and developed in [8], is based on the extrapolation of trajectory $J(t)$ connecting $2^{++}, 4^{++}, 6^{++}$ states (for pomeron) and $3^{--}, 5^{--}, 7^{--}$, states (for odderon) to the physical region of scattering, $t = M^2 \leq 0$. In doing so one assumes that no extra singularities appear in the J plane on the way to $t = 0$. In case of pomeron indeed two other trajectories, f and $f^{(\prime)}$ exist which intersect with glueball trajectory and therefore one should takes into account mixing between them. As a result in [8] a combined trajectory was calculated with realistic pomeron intercept. In this approach perturbative contributions are of subsidiary character and can shift the intercept by approximately 0.2.

¹Note that the usual ω, ρ – Regge poles with $\alpha_u(0) \approx \alpha_\rho(0) \approx 0.5$ give contributions to these reactions, but the corresponding cross sections decrease with energy as $\frac{d\sigma}{dt}|_{t=0} \sim \frac{1}{s}$.

It is important to stress that the region $t \approx 0$ as well as $t \approx M^2 > 0$, $M \sim$ several GeV belongs to the primarily nonperturbative regime, where all intergluonic and interquark distances are large, hence the extrapolation of trajectories to $t = 0$ is inside the NP domain.

In the case of odderon in [8] the intersection of the leading gluonic trajectory with $C = -1$ with $q\bar{q}$ trajectories does not take place, since ρ, ω trajectories have much larger intercepts, and the resulting odderon intercept in [8] obtained using the 3^{--} glueball mass and assumed slope of $(2\pi\sigma_{adj})^{-1}$ was predicted around -1.5. This value excludes possible odderon discovery in reactions $\gamma p \rightarrow (\pi^0, f_2^0, a_2^0)X$ [6, 7] within the upper limits set by the authors.

On the other hand, the perturbative (BFKL) approach [14] starts from another premises. It considers pomeron (odderon) as $2g$ ($3g$) system of Reggeized gluons having only perturbative gluon exchanges, which is justified in the perturbative domain of small ($\lesssim 1 \text{ GeV}^{-1}$) interparticle distances, the situation which might be realized for large negative t , large s , and very small sizes of color dipoles exchanging pomeron. Therefore one encounters the problem of analytic continuation of pomeron singularity to the nonperturbative domain of $t = 0$ (and realistic dipole sizes).

Assuming this can be done, one obtains both pomeron and odderon in vicinity of $J = 1$ [14], see [20], [21] and [5] for more discussion of odderon.

While for pomeron this is reasonable (however the coincidence of pomeron with $J = 1$ is trivial in the limit of small α_s), for odderon it presents a prediction of strong negative C – parity-odd contribution to different reactions, in particular of the type mentioned above [6, 7].

In a different approach(see [4] and refs. therein) a model nonperturbative picture for high-energy scattering was exploited yielding the odderon intercept $\alpha_{odd}(0) = 1$, which as well as the BFKL prediction is at odds with existing data.

Therefore we feel it is necessary to clarify the situation with glueball trajectories and to this end to extend our previous calculations [8] to higher spin states, namely we calculate the masses of glueballs with $C = +1, J = 0^{++}, 2^{++}, 4^{++}, 6^{++}$ and with $C = -1, J = 1^{--}, 2^{--}, 3^{--}, 5^{--}, 7^{--}$. From those we calculate the odderon and pomeron trajectories and find the appropriate intercepts.

Another aim of our calculations is the comparison of newly found glueball masses with existing lattice data [1, 2, 22, 23, 24]. In our previous work [8] we have found a good agreement of all glueball masses with lattice data.

Recently new states, 4^{++} and 6^{++} have been computed [1, 2] and we can compare those with our analytic results. We also compare our results with recent analytic calculations [3, 25, 26]. The paper is organized as follows. In section 2 the Hamiltonian is given together with spin terms following [8] and masses for gg glueballs are calculated.

In section 3 two possible configurations of 3g glueballs are defined, a Δ -type and an Y -type, and the resulting masses are calculated. In section 4 glueball trajectories are obtained and intercepts are found and discussed. The concluding section 5 is devoted to the discussion of the results from the point of view of lattice and experiment correspondence, and to possible improvements.

2 String Hamiltonian and spin corrections

The Hamiltonian for the gg system was derived in [8] in the same way as it was done for the $q\bar{q}$ system [28], and can be written as

$$H = H_0 + \Delta H_s, \quad \Delta H_s = H_{SL} + H_{SS} + H_T \quad (1)$$

where H_0 is the generalization of the spinless $q\bar{q}$ Hamiltonian, obtained in [19], and given in [17, 18]

$$H_0 = \frac{p_r^2}{\mu(t)} + \mu(t) + \frac{L(L+1)}{r^2[\mu + 2\int_0^1(\beta - \frac{1}{2})^2\nu d\beta]} + \\ + \int_0^1 \frac{\sigma_{adj}^2 d\beta}{2\nu(\beta, t)} r^2 + \frac{1}{2} \int_0^1 \nu(\beta, t) d\beta. \quad (2)$$

Here $\mu(t)$ and $\nu(\beta, t)$ are positive auxiliary functions which are to be found from the extremum condition [17]. Their extremal values are equal to the effective gluon energy $\langle\mu\rangle$ and energy density of the adjoint string $\langle\nu\rangle$.

Here $\sigma_{adj} = \frac{9}{4}\sigma_{fund}$, and we shall always set $\sigma_f = 0.18 \text{ GeV}^2$ as found from meson Regge trajectories [17, 29].

To find the spin-averaged masses from (1), $\Delta H_s = 0$, one can use the WKB procedure developed in [30] for the $q\bar{q}$ case and having accuracy better than 5% for $n = 0$, which yields the values given in Table 1 (exact values available for $L = 0$ are given in parentheses).

Table 1

Spin-averaged masses (in GeV) of gg states with $L = 0, \dots, 5$ and $n = 0, 1, 2$, $\sigma_f = 0.18 \text{ GeV}^2$

L n	0	1	2	3	4	5
0	2.09 (2.01)	2.65	3.13	3.53	3.88	4.21
1	3.20 (2.99)	3.65	4.03	4.37	4.67	4.95
2	4.01 (3.75)	4.40	4.74	5.04	5.31	5.56

In (1) and in Table 1 the effect of perturbative gluon exchanges between gluons was neglected, and we consider it as a leading reasonable approximation in obtaining glueball masses. In doing so we follow the argument given in [8], where it was shown that the Coulomb-like adjoint charge interaction is not formed between valence gluons, and moreover, following BFKL approach, one can consider gluon exchange as an effectively not large which can be deduced from the relatively small shift $\Delta \equiv \alpha_P(0) - 1$ of the intercept, when terms $O(\alpha_s^2)$ are taken into account [14, 8]. Therefore in the first approximation we neglect perturbative gluon exchanges, keeping them only in the spin-dependent terms ΔH_S . This strategy is supported by the comparison with lattice data of the spin-averaged masses (see Table 4 of ref. [8]) and, separately, by the comparison of our spin splittings of masses with lattice data, to be discussed below.

Spin-splitting terms ΔH_s are considered in detail in [8] for $L = 0, 1, 2$ and we list in Table 2 the resulting masses, in comparison with existing lattice data.

The difference of Table 2 from the corresponding Table 6 of ref. [8] is that we fix $\sigma_f = 0.18 \text{ GeV}^2$ and recalculate lattice masses for this value of σ_f . Moreover, we take for $L = 0$ in brackets $\alpha_s(ef\bar{f}) = 0.2$ (for spin-splitting terms, since the corresponding spin interaction occurs at relatively small distances, while for the rest masses $\alpha_s(ef\bar{f}) = 0.3$ was taken, as in [8]).

Table 2

Comparison of calculated glueball masses (in GeV) with lattice data ($\sigma_f = 0.18 \text{ GeV}^2$, $\alpha_s = 0.3$ ($\alpha_s = 0.2$ in parentheses))

J^{PC}	M_{theory} this work	M_{lat}		
		[22]	[23]	[24]
0^{++}	(1.61) 1.41	1.53 ± 0.10	1.53 ± 0.04	1.52 ± 0.13
2^{++}	(2.21) 2.30	2.13 ± 0.12	2.20 ± 0.07	2.12 ± 0.15
0^{++*}	(2.72) 2.41	2.38 ± 0.25	2.79 ± 0.09	
2^{++*}	(3.13) 3.32	2.93 ± 0.14	2.85 ± 0.28	
0^{-+}	2.28	2.30 ± 0.15	2.11 ± 0.24	2.27 ± 0.15
0^{-+*}	3.35	3.24 ± 0.2		
2^{-+}	2.70	2.76 ± 0.16	3.0 ± 0.28	2.70 ± 0.19
2^{-+*}	3.73	3.46 ± 0.21		

One can see in Table 2 a good agreement of our calculated glueball masses with measured lattice values, especially for ground states. The radially excited states (marked with an asterix) are some 5-7% higher than lattice data, which probably is the result of WKB approximation [30] for the Hamiltonian (2); indeed as seen from Table 3 of [30], and our Table 1, second column, the exact mass eigenvalues for $L = 0$ and $n_r > 0$ are 6-7% lower than those from WKB approximation.

This agreement of theory with lattice data becomes even more striking, when one realizes that our calculation has no fitting parameters at all, since string tension is a given scale parameter (one could compare dimensionless $M/\sqrt{\sigma}$, as it is done in Table 4 of [8]) and α_s , fixed at the characteristic value, $\alpha_s = 0.3$ describes only spin splitting of masses.

Let us now discuss high spin states, not present in Table 2. The $L = 2, S = 2$ states include $0^{++}, 1^{++}, \dots, 4^{++}$ states which are spread over the mass distance of 63 MeV due to the spin-orbit and tensor level splitting (see table 7 of [8]). Here perturbative and nonperturbative (Thomas term) spin-orbit interaction almost cancel each other.

One can expect, that for higher states the splitting of the states will be even less, as it is observed experimentally for mesons and in what follows we shall neglect spin splitting for states with $L > 2$.

The resulting glueball masses are given in Table 3.

Table 3

Glueball masses (in GeV) for $L \geq 2$, from the Hamiltonian (2)

L, S	States	Masses
2,2	$0^{++}, 4^{++}, 1^{++}, 2^{++}, 3^{++}$	3.11-3.17
2,0	2^{++}	3.13
3,1	$4^{-+}, 3^{-+}, 2^{-+}$	3.53
4,2	$6^{++} - 2^{++}$	3.88
4,0	4^{++}	3.88
5,1	$6^{-+}, 5^{-+}, 4^{-+}$	4.21

These masses can be compared to (scarce) lattice values. E.g. $M_{lat}(3^{++}) = 3.28 \pm 0.2$ GeV in [22], 3.81 ± 0.31 in [23] and 3.26 ± 0.21 in [24]; $M_{lat}(4^{++}) = 3.65$ GeV in [2] and around 3.5 GeV in [1]; $M_{lat}(6^{++}) = 4.2 \pm 0.2$ GeV in [1].

One can notice an approximate degeneracy of 3^{++} and 4^{++} states on the lattice in agreement with theory. For the pomeron trajectory the state $6^{++}(L=4, S=2)$ is important and will be used below.

It follows from Table 3, that there are several states with the same J^P which do not differ substantially in mass (for example 4^{++} in (2,2), (4,2) and (4,0) configurations). This makes difficult to compare predicted masses for these states with lattice data, which are usually approximated by a single state. Note that masses of states with $J^P = 4^+6^+$ on the pomeron trajectory calculated in ref [1] are higher than the lowest values, obtained in our paper. As a result the slope of the pomeron trajectory obtained in paper [1] is smaller than the value corresponding to Casimir scaling and an intercept of the purely glueball pomeron is higher than in the present approach. As we discuss in Section 4 below this intercept has little to do with a physical pomeron intercept due to large mixing with $q\bar{q}$ trajectories.

3 Three-gluon glueballs

The $3g$ glueballs (oddballs) can be of two basic configurations: the Δ -type and the Y -type (only the first one was considered in [8]). The corresponding wave operators for the Δ -type are given in Table 10 of [8] and basically

correspond to 3 gluons sitting in vertices of a triangle and connected by the fundamental strings. Another, not considered in [8] the Y , form is composed as

$$\Psi(x_1, x_2, x_3) \sim d_{abc} E_i^a(x) E_k^b(x) E_l^c(x) \quad (3)$$

and is the adjoint equivalent of the baryon operator with the replacement $e_{\alpha\beta\gamma} \rightarrow d_{abc}$. Another possible Y -type form is made with the operator f_{abc} instead of d_{abc} and requires antisymmetric spin-coordinate function. Using the charge-conjugation C transformation $F_{\mu\nu} \rightarrow -F_{\mu\nu}^T$, one can easily understand that the form made of f_{abc} has $C = +1$. Both f, d forms have been used in [3].

The spin-independent part of Hamiltonian in both cases can be written as

$$H_0^{(3g)} = T_{3g} + \frac{3\mu}{2} + V_{Y,\Delta}(\mathbf{r}_1, \mathbf{r}_2, \mathbf{r}_3) \quad (4)$$

where T_{3g} is the kinetic operator,

$$T_{3g} = \frac{\mathbf{p}_\eta^2 + \mathbf{p}_\xi^2}{2\mu}, \quad \mathbf{p} = \frac{1}{i} \frac{\partial}{\partial \xi}, \quad \mathbf{p}_\eta = \frac{1}{i} \frac{\partial}{\partial \eta} \quad (5)$$

and the Jacobi coordinates defined as

$$\xi = \sqrt{\frac{3}{2}} \left(\frac{\mathbf{r}_1 + \mathbf{r}_2}{2} - \mathbf{r}_3 \right), \quad \eta = \frac{\mathbf{r}_1 - \mathbf{r}_2}{\sqrt{2}}. \quad (6)$$

Finally the interaction is

$$V_\Delta = \sigma_f \sum_{i < j} |\mathbf{r}_i - \mathbf{r}_j|, \quad V_Y = \sigma_{adj} \sum_{i=1}^3 |\mathbf{r}_i - \mathbf{R}_Y| \quad (7)$$

and $\mathbf{R}_Y$ is the position of the string junction. We shall be using $\sigma_f \equiv \sigma$, $\sigma_{adj} = \frac{C_2(adj)}{C_2(fund)} \sigma = \frac{9}{4} \sigma$.

We are solving equation $H_0^{(3g)} \Psi = M(\mu) \Psi$ as in [8] using the hyperspherical approach [31, 32], which yields very good accuracy already in the lowest approximation [33]. Defining the hyperradius $\rho, \rho^2 = \eta^2 + \xi^2$, and grand orbital momentum K , $K = L, L + 2, L + 4, \dots$, one has the equation

$$-\frac{1}{2\mu} \frac{d^2 \chi}{d\rho^2} + U_{\Delta,Y}(\rho) \chi(\rho) = \varepsilon(\mu) \chi(\rho), \quad (8)$$

with

$$U_{\Delta,Y}(\rho) = \frac{1}{2\mu\rho^2} \left(K^2 + 4K + \frac{15}{4} \right) + c_{\Delta,Y}\rho\sigma, \quad (9)$$

$$c_{\Delta} = \frac{32\sqrt{2}}{5\pi} = 2.88; \quad c_Y = 3.31. \quad (10)$$

Here one notice that the case of Y form can be obtained from baryonic calculations of [32] by a simple replacement $\sigma_f \rightarrow \sigma_{adj}$ (see [32] for details of derivation) and the total mass $M(3g)$ is obtained by minimizing the mass $M(\mu)$ over the values of μ ,

$$M(3g) = \min_{\mu} M(\mu) = \min_{\mu} \left\{ \frac{3\mu}{2} + \varepsilon(\mu) \right\}. \quad (11)$$

It was shown [33], that $\varepsilon(\mu)$ can be found (with one percent accuracy) from the minimum of $U_{\Delta,Y}(\rho)$ at some point $\rho = \rho_0$.

In this way one obtains for $\varepsilon(\mu)$.

$$\varepsilon_{\Delta,Y}(\mu) = \frac{3}{2} \frac{(c_{\Delta,Y})^{2/3} \sigma^{2/3}}{\mu^{1/3}} k^{1/3} \left(1 + \frac{1}{\sqrt{3k}} \right) \equiv \frac{\Omega_{\Delta,Y} \sigma^{2/3}}{\mu^{1/3}} \quad (12)$$

where $k \equiv K^2 + 4K + \frac{15}{4}$.

Minimizing over μ in (11), one finds the constituent gluon mass μ_0 ,

$$\mu_0 = \left(\frac{2}{9} \Omega_{\Delta,Y} \right)^{3/4} \sqrt{\sigma}, \quad M(3g) = 6\mu_0. \quad (13)$$

One can now predict the $3g$ glueball masses, still without spin splittings, fixing the value of $K_{min} = L = 0, 1, 2, \dots$

For the lowest $3g$ state with $K_{min} = L = 0$, the spin splitting is due to the hyperfine interaction and it was calculated in [8], yielding

$$\Delta M_{ss} = 0.644\mu_0 \frac{J(J+1) - 6}{6} \quad (14)$$

This gives $\Delta M_{ss}(3^{--}) = 0.28 \text{ GeV}$, $\Delta M_{ss}(2^{--}) = 0$, $\Delta M_{ss}(1^{--}) = -0.189 \text{ GeV}$.

For $L > 0$ splitting is due to tensor and spin-orbit forces, and we assume, that the situation is similar to that of baryons, where these forces are known to be weak. The situation might however be different for $3g$ states.

We neglect spin splittings for $L > 0$ and give in Table 4 the calculated spin-averaged masses. One should note that the Y -type glueballs are fully equivalent to baryons and their masses are obtained by simply multiplying baryon masses from [32] by the factor $\sqrt{\frac{C_2(adj)}{C_2(fund)}} = \frac{3}{2}$. More sophisticated configurations were also considered in [32] however the resulting masses are very close to the corresponding M_Y values and are omitted from the Table 4.

Table 4

Oddball masses (in GeV) for various $L = 0, 1, \dots, 4$ and $J \leq 7$ in comparison with lattice results. For $L > 0$ only spin-averaged values are shown, $\sigma = 0.18 \text{ GeV}^2$.

L	J^{PC}	$M_\Delta(theor)$ this work	$M_Y(theor)$ this work	$M(lat)$ [24]	[1]
0	1^{--}	3.02	3.32	3.40 ± 0.21	3.10
	2^{--}	3.21	3.53	3.56 ± 0.21	3.55
	3^{--}	3.49	3.83	3.73 ± 0.21	4.15
1	$0^{+-}, \dots, 3^{+-}$	3.72	4.09	3.46 ± 0.2	~ 3.2
2	$5^{--}, \dots, 1^{--}$	4.18	4.59		
4	$7^{--}, \dots, 1^{--}$	4.96	5.25		

One can see from Table 4, that agreement between theoretical and lattice values is reasonable and of the same quality, as the agreement between results of different lattice groups. Our results are closer to the lattice data of [22, 24] and lie below those of [1]; the spin splittings in the $(1^{--}, 2^{--}, 3^{--})$ triplet are 0.47 GeV in our calculation [8] and 0.33 ± 0.21 in data of [22, 24], which again may indicate that effective α_s for spin-spin interaction is $\alpha_s = 0.2$ rather than our fixed value $\alpha_s = 0.3$.

Two other models have been used in [3] for $1^{--}, 2^{--}, 3^{--}, 5^{--}, 7^{--}$ states with results similar to ours for the first triplet of states, but much heavier for $5^{--}, 7^{--}$ states.

4 Pomeron and odderon trajectories

Since our results are the same as in [8] for $L = 0, 2$, and the resulting pomeron trajectory is assumed to be the same as in [8] (mixed with f, f' trajectories), it is worthwhile to compare the slopes $\alpha'_p(0)$ obtained from the standard procedure, $\alpha'_p(stand) = \frac{1}{2\pi\sigma_{adj}} = 0.393 \text{ GeV}^{-2}$, with the slopes obtained from masses $M(J = 2^{++}; 4^{++}; 6^{++}) = (2.3 - 2.21; 3.15; 3.88) \text{ GeV}$, where we give the 2^{++} mass interval for $\alpha_s = 0.3 - 0.2$.

One obtains for the neighboring masses

$$\alpha'_p(4^{++} - 2^{++}) = (0.431 \div 0.397) \text{ GeV}^2, \quad \alpha'_p(6^{++} - 4^{++}) = 0.3971 \text{ GeV}^2. \quad (15)$$

Thus we see that the slopes are close to the standard one, and the trajectory is close to the straight line. The value of intercept, on the other hand depends crucially on the intersection with f, f' trajectories and as we argued in [8], this value is not actually controlled by the masses on the pomeron trajectory. This is in contrast to the case of odderon, where no intersection with meson trajectories is possible for $t > 0$ and hence intercept can be estimated from the computed above masses, and now we have three states on the trajectory and can determine both slope and intercept.

We define two odderon trajectories corresponding to the Δ -type and Y -type configurations,

$$\alpha_\Delta(t) = \alpha'_\Delta \cdot t + \alpha_\Delta(0), \quad \alpha_Y(t) = \alpha'_Y \cdot t + \alpha_Y(0). \quad (16)$$

It appears that one can find odderon trajectory analytically, using explicit mass formulas Eqs. (12)-(14). Here spin splittings are neglected, as it usually done for meson Regge trajectories [29]. E.G. in the latter case spin dependent correlations split the spin-averaged Regge trajectory into the two nonlinear trajectories with $J = L \pm 1$.

In this way one can approximate k in (13) for $K = L$, as $k = (L+2)^2 - \frac{1}{4} \approx (L+2)^2$ and hence

$$M_\Delta^2(3g)/\sigma = 4c_\Delta \sqrt{3k} \left(1 + \frac{1}{\sqrt{3k}}\right)^{3/2} \approx 20.0 \left(L + 2 + \frac{\sqrt{3}}{2}\right). \quad (17)$$

Taking into account, that $J_{odd} = L+3$, one obtains the odderon trajectory

$$M^2(3g) = 20.0\sigma(J - \alpha_\Delta(0)), \quad \alpha_\Delta(0) = 1 - \frac{\sqrt{3}}{2} = 0.134. \quad (18)$$

At this point one can compare oddball masses from (18) for $J = 3^{--}, 5^{--}$ and 7^{--} with the values given in the Table 4, and find that these masses differ less than 1%, if for $M(3^{--})$ one takes the spin-averaged value, almost coinciding with $M(2^{--}) = 3.21$.

It is interesting to compare the resulting slope in (18) $\alpha'_\Delta = \frac{1}{20\sigma} = 0.277 \text{ GeV}^{-2}$ with the standard relativistic "potential" slope (when string rotation correction is not taken into account which is our case), $\alpha'_\Delta(pot) = \frac{1}{8\sigma_a} = 0.308 \text{ GeV}^{-2}$. The string correction was computed in [17, 29] and it decreases the glueball masses with $L \neq 0$ by $4 \div 5\%$ and the asymptotic ($L \gg 1$) slope is $\alpha'_\Delta(string) = \frac{1}{2\pi\sigma_a} = 0.39 \text{ GeV}^{-2}$. If one uses this slope (and keeps the mass with $L = 0$ unchanged), then one has the string-corrected odderon trajectory

$$M_{string}^2(3g) = 2\pi\sigma_a(J - \alpha_\Delta^{string}(0)), \quad \alpha_\Delta^{string}(0) = -1.05. \quad (19)$$

It is clear from (11), (13), that masses of Y -type glueballs are higher, namely $\frac{M_Y}{M_\Delta} = \sqrt{\frac{c_Y}{c_\Delta}} \cong 1.1$, and the intercept of Y trajectory is lower.

The same procedure as for Eq. (19) leads for the intercept of Y -type trajectory the value

$$\alpha_Y^{string}(0) = -1.9. \quad (20)$$

Now if one insists on using spin-splitted (nonlinear in general) trajectory passing through 3^{--} and 5^{--} states, one obtains extrapolating $\alpha_\Delta(t)$ to $t = 0$ the intercept $\alpha_\Delta = -1.6$.

We see that determination of intercepts of $3g$ -trajectories is model dependent, but in both cases, Eq. (18) and Eq. (19), the odderon intercept is far from unity.

5 Discussion and summary

As shown in the previous section, the odderon intercept is 0.14 (without string corrections) and drops down to -1.05 when the string slope $\frac{1}{2\pi\sigma_a}$ is restored. In any case intercepts of these $3g$ -trajectories are very far from the perturbative intercept $\alpha_{3g} \approx 1$. Thus the nonperturbative effects strongly reduce the intercept of the $3g$ -trajectories and there is no "odderon" (the singularity with negative $C = \sigma$ and $\alpha_o(0) \approx 1$) in our approach.

This difference for the intercepts of $3g$ -singularities with results of perturbation theory is of principle importance and takes place for all multigluonic states. In the perturbation theory the multigluon singularities in j -plane are

above unity [34] and in principle mix with $2g$ -state. Account of nonperturbative interaction between gluons lead to large masses of multigluon states and consequently to low intercepts of such trajectories. Qualitatively it can be understood as follows: NP interactions confine and create effective mass for each gluon (calculable in FCM [15, 16]) proportional to $\sqrt{\sigma}$. Therefore $4g$ states are separated from $2g$ states by an interval ~ 2 GeV and mixing can be neglected in the first approximation. This is contrast to BFKL method, where such interval is absent and in principle all multigluon states should be considered simultaneously.

Comparison of our calculated masses with lattice data in Tables 2,3 shows that the ordering of states in mass values is the same. The agreement in masses for states with $L = 0, 1$ and $n_r = 0$ is surprisingly good although no fitting parameters are used. The (small) discrepancy for the states with $n_r > 0$ was discussed above, and it is planned to improve theoretical accuracy for $L > 0, n_r > 0$. For $L \geq 2$ our spin-averaged values are in a good agreement with lattice data of [22, 24] and 10÷20% below data of [1, 2, 23]. One should take into account at this point, that 4^{++} and 6^{++} states occur from $L = 2$ and 4, and $L = 4$ and 6 respectively, and resulting mixing can shift the masses from the one-channel values in lattice computations (as well as in our analytic approach).

For $3g$ spin-averaged states in Table 3 the agreement with lattice data from [24, 1] is within 10%.

Our results for the odderon masses of $J^{PC} = 5^{--}, 7^{--}$ are some 0.5 GeV below the results of models H_{eff}^g and H_M from [3], however qualitatively the odderon intercepts are also low (-0.88 and 0.25 respectively).

In summary, we have calculated high spin states of both $2g$ and $3g$ systems and found the corresponding masses, in this way extending our results from an earlier paper [8]. We have confirmed our previous results on the pomeron trajectory.

In the case of odderon we have found two trajectories, of Δ and Y type with masses differing by $\sim 10\%$ and similar intercepts below 0.14. This result might explain why odderon is not seen in photonucleon reactions [6, 7], and is in sharp contrast to the BFKL-type odderon intercept, which is around 1. This work is supported by the Federal Program of the Russian Ministry of industry, Science and Technology No.40.052.1.1.1112, and by the grant for scientific schools NS-1774. 2003. 2. One of authors (A.B.K.) acknowledges partial support of the grants CRDF RUP2-2621-M0-04 and 04-02-17263.

References

- [1] H.B.Meyer, M.J.Teper, Phys. Lett. **B605** (2005) 344.
- [2] Da Quing Liu, Ji Min Wu, hep-lat/0105019.
- [3] F.J.Llanes-Estrada P.Bicudo, S.R.Cotanch, hep-ph/0507205.
- [4] A. Donnachie, H.G.Dosch, O.Nachtmann, hep-ph/0508196.
- [5] C.Ewerz, hep-ph/0511196; hep-ph/0306137.
- [6] C.Adloff et al. H1 Collaboration, Phys. Lett. **B544** (2002) 35.
- [7] T.Berndt (for the H1 Collaboration), Acta Phys. Polonica **B33** (2002) 3499 .
- [8] A.B.Kaidalov, Yu.A.Simonov, Phys. Lett. **B 477** (2000) 163; hep-ph/9912434; Phys. At. Nucl. **63** (2000) 1428, hep-ph/9911291.
- [9] H.Fritzsch and P.Minkowski, Nuovo Cim. **A30** (1975) 393; M.Fritzsch and M.Gell–Mann, Proc. 16th Int. Conf. on High-Energy Physics, FNAL 1972 V.2. P. 135.
- [10] R.L.Jaffe and K.Johnson, Phys. Lett. **B60** (1976) 201.
- [11] P.Hasenfrantz and J.Kuti, Phys. Rep. **40** (1978) 75.
- [12] D.Robson Nucl. Phys. **B130** (1977) 328;
J.M.Cornwall and A.Soni, Phys. Lett. **B120** (1983) 431.
- [13] F.E.Low, Phys. Rev. **D12** (1975) 163;
S.Nussinov, Phys. Rev. Lett. **34** (1975) 1286.
- [14] V.S.Fadin, E.A.Kuraev, L.N.Lipatov, Sov. Phys. JETP **44** (1976) 443;
45 (1977) 199; I.I.Balitsky, L.N.Lipatov, Sov. J. Nucl. Phys. **28** (1978) 822;
L.N.Lipatov, Nucl. Phys. **B365** (1991) 614; Sov. Phys. JETP **63** (1986) 904;
J.Bartels, Nucl. Phys. **B175** (1980) 365; J.Kwiecinski, M.Praszalowicz, Phys. Lett. **B94** (1980) 413.
- [15] H.G.Dosch, Phys. Lett. B **190** (1987) 177;
H.G.Dosch and Yu.A.Simonov, Phys. Lett. B **205** (1988) 339;
Yu.A.Simonov, Nucl. Phys. B **307** (1988) 512.

- [16] A.Di Giacomo, H.G.Dosch, V.I.Shevchenko, Yu.A.Simonov, Phys. Rept. **372** (2002) 319; hep-ph/0007223.
- [17] A.Yu.Dubin, A.B.Kaidalov, and Yu.A.Simonov, Phys. Lett. **B323** (1994) 41; Yad. Fiz. **56** (1993) 213.
- [18] Dan La Course and M.G.Olsson, Phys. Rev. **D39** (1989) 2751; M.G.Olsson, Nuovo Cim., **107 A** (1994) 2541.
- [19] Yu. A.Simonov, Phys. Lett. **B 249** (1990) 514.
- [20] L.Lukaszuk, B.Nicolescu, Lett. Nuovo. Cim **8** (1973) 405.
- [21] B.Nicolescu, Nucl. Phys. Proc. Suppl. **25B** (1992) 142; hep-ph/9911334.
- [22] C.Morningstar, M.Peardon, Nucl. Phys. **B63** (1998) 22; (Proc. Suppl.); Phys. Rev. **D60** (1999) 034509.
- [23] M. Teper, hep-th/98121
- [24] Y.Chen et al., hep-lat/0510074
- [25] F.Brau, C.Semay, hep-ph/0412173.
- [26] V.Mathieu, C.Semay, F.Brau, hep-ph/0511210.
- [27] A.P.Szczepaniak, E.S.Swanson, Phys. Lett. **B577** (2003) 61.
- [28] Yu.A.Simonov, in: "QCD: Perturbative or nonperturbative?" L.S.Ferreira, P.Noguera, and J.I.Silva-Marcos eds. Proceedings of the XVII Quantum School, Lisbon, 29 Sept. – 4 Oct., 1999, p. 60 (World Sci., 2000).
- [29] A.M.Badalian, B.L.G. Bakker, Phys. Rev. **D66** (2002) 034025, hep-ph/0202246.
- [30] V.L.Morgunov, A.V.Nefediev, and Yu.A.Simonov, Phys. Lett. **B459** (1999) 653; hep-ph/9906318.
- [31] Yu.A.Simonov, Yad.Fiz., **3** (1966) 630; (Sov.J. Nucl. Phys., **3** (1966) 461; A.M.Badalian and Yu.A.Simonov, Yad.Fiz. **3** (1966) 1032; **5** (1967) 88; (Sov. J. Nucl. Phys. **3** (1966) 755; **5** (1967) 60; F.Calogero and Yu.A. Simonov, Phys. Rev., **169** (1968) 789.

- [32] Yu.A.Simonov, Phys. Lett. **B228** (1989) 413; M.Fabre de la Ripelle and Yu. A.Simonov, Ann. Phys. (N.Y.) **212** (1991) 235.
- [33] B.O.Kerbikov, M.I.Polikarpov, L.V.Shevchenko, Nucl. Phys. **B331** (1990) 19;
I.M.Narodetskii, M.A.Trusov, Phys. Atom. Nucl. **67** (2004) 762.
- [34] H.J. de Vega, L.N.Lipatov, Phys. Rev. **D66** (2002) 074013;
G.P.Korchemsky, J.Kotanski, A.N.Manashov, Phys. Lett. **B583** (2004)121; Phys. Rev. Lett. **88** (2002) 122002.